

An axiomatic approach in minimum cost spanning tree problems with groups

Gustavo Bergantiños, Research Group in Economic Analysis
Universidade de Vigo (Spain).*

María Gómez-Rúa, Research Group in Economic Analysis
Universidade de Vigo (Spain)

Abstract

We study minimum cost spanning tree problems with groups, where agents are located in different villages, cities, etc. The groups are the agents of the same village. In Bergantiños and Gómez-Rúa (2009, *Economic Theory*) we define the rule F as the Owen value of the irreducible game with groups and we prove that F generalizes the folk rule of minimum cost spanning tree problems. Bergantiños and Vidal-Puga (2007, *Journal of Economic Theory*) give two characterizations of the folk rule. In the first one they characterize it as the unique rule satisfying solidarity, population monotonicity and equal share of extra costs. In the second characterization of the folk rule they replace solidarity by independence of irrelevant trees and population monotonicity by separability. In this paper we extend such characterizations to our setting. Some of the properties are the same (solidarity and independence of irrelevant trees) and the other need to be adapted. In general, we do it by claiming the property twice: once among the groups and the other among the agents inside the same group.

Keywords: minimum cost spanning tree problems, folk rule, cost sharing, axiomatization.

*Corresponding author: Tel: +34986812497; fax: +34986812401. E-mail: gbergant@uvigo.es

1 Introduction

In minimum cost spanning tree problems, *mcstp*, a group of agents want to be connected to a single supplier of some service. Usually, agents can reduce the total cost if some agents connect to the supplier through other agents. The cheapest graph connecting all agents to the supplier is called the minimum cost spanning tree, *mt*. It is assumed that agents construct an *mt*. Now a cost allocation problem arises. The most relevant question is how to divide the cost of the *mt* among the agents.

The most popular way for answering this question is through the axiomatic characterization of a single rule (see for instance, Kar (2002), Dutta and Kar (2004), and Branzei *et al* (2004)) or a family of rules (see for instance, Bergantiños *et al* (2009), Bogomolnaia and Moulin (2008), and Chun and Lee (2009)).

Sometimes agents are located in different villages (see for instance, Dutta and Kar (2004) or Bergantiños and Lorenzo (2004, 2005, 2008)). This means, in terms of the cost matrix, that the connection cost between two agents of the same village is not larger than the connection cost between an agent of this village and an agent from other village. The classical model of *mcstp* includes these situations. Nevertheless, it ignores the fact that some group of agents are located in the same village. In Bergantiños and Gómez-Rúa (2009) we introduce a new class of problems which includes this fact in the model. We do it by considering an extra element in the model, namely a partition $G = \{G^1, \dots, G^m\}$ of the set of agents N . For each $k = 1, \dots, m$, G^k represents the group of agents located in the same village, city, ... We call these problems *mcstp* with groups.

A very well-known rule in *mcstp* is the folk rule. It was introduced in Feltkamp *et al.* (1994) and studied later in further papers, for instance Branzei *et al.* (2004) and Bergantiños and Vidal-Puga (2007a, 2007b, 2009). The folk rule can be defined in different ways, for instance, through Kruskal's algorithm or as the Shapley value of the irreducible game or the optimistic game. In Bergantiños and Gómez-Rúa (2009) we define the rule F in *mcstp* with groups as the Owen value (Owen, 1977) of the irreducible game with groups. Since the Owen value generalizes the Shapley value, F coincides with the folk rule in classical *mcstp*.

Bergantiños and Vidal-Puga (2009) give an axiomatic characterization of the folk rule. In Bergantiños and Gómez-Rúa (2009) we extend it to *mcstp* with groups. Some of the properties are the same and other need to be

adapted. We do it by claiming the property twice: once among the groups and the other among the agents inside the same group. Let us clarify this idea with the property of population monotonicity (PM). In classical $mcstp$ PM says that if a new agent joins the society, no agent of the initial society can be worse off. In $mcstp$ with groups PM is divided in two parts: Population monotonicity over groups (PMG) and population monotonicity over agents (PMA). PMG says that if a new group joins the society, no agent of the initial society can be worse off. PMA says that if agent i enters in group G^k , no agent of group G^k can be worse off. Moreover, if the connection costs between group G^k and the other groups do not change, agents of the other groups must pay the same.

Bergantiños and Vidal-Puga (2007a) give two characterizations of the folk rule. In the first one they prove that the folk rule is the unique rule satisfying solidarity (SOL), PM , and equal share of extra costs ($ESEC$). SOL states that if a network connection cost increases, no agent should pay less. The idea behind $ESEC$ is the following: consider a problem where the most expensive connection cost for any agent is the cost of connecting to the source. Moreover, the connection cost to the source is the same for all agents. If we assume that this connection cost increases by $x > 0$, then $ESEC$ states that if agent i pays f_i in the original problem, he must pay $f_i + \frac{x}{n}$ when the cost increases (where n is the set of agents). In the second characterization of the folk rule they replace SOL by independence of irrelevant trees (IIT) and PM by separability (SEP). IIT states that if two $mcstp$ have the same mt with the same costs in each arc belonging to, then the rule must coincide in both $mcstp$. Assuming that two subsets of agents, S and $N \setminus S$, can be connected to the source either separately or jointly, SEP states that if there are no savings when they are jointly connected to the source, then agents will pay the same in both circumstances.

In this paper we present two characterizations of the rule F in $mcstp$ with groups, generalizing the results of Bergantiños and Vidal-Puga (2007a). We first adapt the properties to $mcstp$ with groups. SOL and IIT do not need to be adapted (both properties make sense when there are groups). $ESEC$ is replaced by coalitional sharing of extra costs ($CSEC$). Under the hypothesis of $ESEC$, $CSEC$ states that the groups should share this extra cost x equally among them. Moreover, the extra cost assigned to each group should be shared equally among the agents of this group. PM is divided in two properties PMG and PMA , as in Bergantiños and Gómez-Rúa (2009). Finally, SEP is also divided in two properties: separability among groups

(*SEPG*) and separability among agents (*SEPA*).

The results establish that F is the unique rule in *mcstp* with groups satisfying *SOL*, *PMA*, *PMG*, and *CSEC*. We also prove that F is the unique rule in *mcstp* with groups satisfying *IIT*, *SEPA*, *SEPG*, and *CSEC*; and finally we obtain that the properties used in both results are independent.

The paper is organized as follows. In Section 2 we introduce *mcstp*. In Section 3 we introduce *mcstp* with groups and several properties. In Section 4 we present our results.

2 The Problem

Let $\mathcal{N} = \{1, 2, \dots\}$ be the set of all possible agents. Given a finite set $N \subset \mathcal{N}$, let Π_N be the set of all permutations over N . Given $\pi \in \Pi_N$, let $Pre(i, \pi)$ denote the set of elements of N which come before i in the order given by π , *i.e.*, $Pre(i, \pi) = \{j \in N : \pi(j) < \pi(i)\}$.

We are interested in networks whose nodes are elements of a set $N_0 = N \cup \{0\}$, where $N \subset \mathcal{N}$ is finite and 0 is a special node called the *source*. Usually we take $N = \{1, \dots, n\}$.

A *cost matrix* $C = (c_{ij})_{i,j \in N_0}$ on N represents the cost of direct link between any pair of nodes. We assume that $c_{ij} = c_{ji} \geq 0$ for each $i, j \in N_0$ and $c_{ii} = 0$ for each $i \in N_0$. Since $c_{ij} = c_{ji}$ we work with undirected arcs, *i.e.* $(i, j) = (j, i)$.

We denote the set of all cost matrices over N as \mathcal{C}^N . Given $C, C' \in \mathcal{C}^N$ we say $C \leq C'$ if $c_{ij} \leq c'_{ij}$ for all $i, j \in N_0$.

A *minimum cost spanning tree problem*, briefly an *mcstp*, is a pair (N_0, C) where $N \subset \mathcal{N}$ is a finite set of agents, 0 is the source, and $C \in \mathcal{C}^N$ is the cost matrix.

Given an *mcstp* (N_0, C) , we define the *mcstp* induced by C in $S \subset N$ as (S_0, C) .

A *network* g over N_0 is a subset of $\{(i, j) : i, j \in N_0, i \neq j\}$. The elements of g are called *arcs*. Given a network g over N_0 and $S \subset N_0$ we denote by g_S the network induced by g among the elements of S . Namely, $g_S = \{(i, j) \in g : \{i, j\} \subset S\}$.

Given a network g and a pair of nodes i and j , a *path* from i to j in g is a sequence of different arcs $\{(i_{h-1}, i_h)\}_{h=1}^l$ satisfying $(i_{h-1}, i_h) \in g$ for all $h \in \{1, 2, \dots, l\}$, $i = i_0$, and $j = i_l$.

A *tree* is a network such that for all $i \in N$ there is a unique path from i to

the source. If t is a tree, we usually write $t = \{(i^0, i)\}_{i \in N}$ where i^0 represents the first agent in the unique path in t from i to 0.

Let \mathcal{G}^N denote the set of all networks over N_0 . Let \mathcal{G}_0^N denote the set of all networks where every agent $i \in N$ is connected to the source, *i.e.* there exists a path from i to 0 in the network.

Given an *mcstp* (N_0, C) and $g \in \mathcal{G}^N$, we define the *cost* associated with g as

$$c(N_0, C, g) = \sum_{(i,j) \in g} c_{ij}.$$

When there is no ambiguity, we write $c(g)$ or $c(C, g)$ instead of $c(N_0, C, g)$.

An *minimum cost spanning tree* for (N_0, C) , briefly an *mt*, is a tree t over N_0 such that $c(t) = \min_{g \in \mathcal{G}_0^N} c(g)$. It is well-known that an *mt* exists, even though it is not necessarily unique. Given an *mcstp* (N_0, C) , we denote the cost associated with any *mt* as $m(N_0, C)$.

Given an *mcstp*, Prim (1957) provides an algorithm for solving the problem of connecting all agents to the source such that the total cost of creating the network is minimal. The idea of this algorithm is simple: starting from the source we construct a network by sequentially adding arcs with the lowest cost and without introducing cycles.

Formally, Prim's algorithm is defined as follows. We start with $S^0 = \{0\}$ and $g^0 = \emptyset$.

Stage 1: Take an arc $(0, i_1)$ such that $c_{0i_1} = \min_{j \in N} \{c_{0j}\}$. If there are several arcs satisfying this condition, select just one. Now, $S^1 = \{0, i_1\}$ and $g^1 = \{(0, i_1)\}$.

Stage $p + 1$: Assume that we have defined $S^p \subset N_0$ and $g^p \in \mathcal{G}^N$. We now define S^{p+1} and g^{p+1} . Take an arc (i_{p+1}^0, i_{p+1}) with $i_{p+1}^0 \in S^p$ and $i_{p+1} \in N_0 \setminus S^p$ such that $c_{i_{p+1}^0 i_{p+1}} = \min_{k \in S^p, l \in N_0 \setminus S^p} \{c_{kl}\}$. If there are several arcs satisfying this condition, select just one. Now, $S^{p+1} = S^p \cup \{i_{p+1}\}$ and $g^{p+1} = g^p \cup \{(i_{p+1}^0, i_{p+1})\}$.

This process is completed in n stages. We say that g^n is a tree obtained following Prim's algorithm. Notice that this algorithm leads to a tree, but this is not always unique.

Given a tree t , we define the *predecessor set* of a node i in t as $Pre(i, t) = \{j \in N_0 : j \text{ is in the unique path from } i \text{ to the source}\}$. We assume that $i \notin Pre(i, t)$ and $0 \in Pre(i, t)$ when $i \neq 0$. For notational convenience, $Pre(0, t) = \emptyset$. The *distance* from node i to the source in t is the cardinality

of $Pre(i, t)$. The *immediate predecessor* of agent i in t , denoted as i^0 , is the node that comes immediately before i , that is, $i^0 \in Pre(i, t)$ and $k \in Pre(i, t)$ implies either $k = i^0$ or $k \in Pre(i^0, t)$. Note that $Pre(i^0, t) \subset Pre(i, t)$ and $Pre(i, t) \setminus Pre(i^0, t) = \{i^0\}$. The *follower set* of an agent i in t is the set $Fol(i, t) = \{j \in N : i \in Pre(j, t)\}$.

Given an *mcstp* (N_0, C) and an *mt* t , Bird (1976) defined the *minimal network* (N_0, C^t) associated with t as follows: $c_{ij}^t = \max_{(k,l) \in g_{ij}} \{c_{kl}\}$, where g_{ij} denotes the unique path in t from i to j . Even though g_{ij} depends on the choice of t , c_{ij}^t is independent of the chosen t . Proof of this can be found, for instance, in Aarts and Driessen (1993).

The *irreducible form* of an *mcstp* (N_0, C) is defined as the minimal network (N_0, C^*) associated with a particular *mt* t . If (N_0, C^*) is an *irreducible form*, we say that C^* is an *irreducible matrix*.

A *coalitional game with transferable utility*, briefly a *TU game*, is a pair (N, v) where $v : 2^N \rightarrow \mathbb{R}$ satisfies $v(\emptyset) = 0$. $Sh(N, v)$ denotes the Shapley value (Shapley, 1953) of (N, v) .

For each *mcstp* (N_0, C) . Bird (1976) introduces the *TU game* (N, v_C) . For each coalition $S \subset N$,

$$v_C(S) = m(S_0, C).$$

One of the most important issues addressed in the literature about *mcstp* is how to divide the cost of connecting agents to the source between them. We now briefly introduce some of the rules studied in the literature.

A (*cost allocation*) *rule* is a function f such that $f(N_0, C) \in \mathbb{R}^N$ and $\sum_{i \in N} f_i(N_0, C) = m(N_0, C)$ for each *mcstp* (N_0, C) . As usual, $f_i(N_0, C)$ represents the cost allocated to agent i .

Notice that we implicitly assume that the agents build an *mt*. As far as we know, all the rules proposed in the literature make this assumption.

There are several rules studied in the literature. We mention, for instance, the rules studied in Bird (1976), Kar (2002), and Dutta and Kar (2004). In this paper the rule introduced by Feltkamp *et al.* (1994) and called Equal Remaining Obligations rule (*ERO*) will be very important. *ERO* is called the *P - value* in Branzei *et al.* (2004).

On the other hand, in Bergantiños and Vidal-Puga (2007a) the rule φ is defined as

$$\varphi(N_0, C) = Sh(N, v_{C^*})$$

where C^* is the irreducible matrix associated with C . Surprisingly φ coincides with ERO , a proof can be found, for instance, in Bergantiños and Lorenzo-Freire (2008). This rule is also studied in Bergantiños and Vidal-Puga (2007b, 2009). We call *folk rule* to this rule.

We now present some properties introduced in the literature of the *mcstp*.

Population Monotonicity (PM): For all *mcstp* (N_0, C) , $S \subset N$, and $i \in S$, we have

$$\psi_i(N_0, C) \leq \psi_i(S_0, C).$$

This property implies that if new agents join a "society" no agent from the "initial society" can be worse off.

Separability (SEP): For all *mcstp* (N_0, C) and $S \subset N$ satisfying $m(N_0, C) = m(S_0, C) + m((N \setminus S)_0, C)$, we have

$$f_i(N_0, C) = \begin{cases} f_i(S_0, C) & \text{if } i \in S \\ f_i((N \setminus S)_0, C) & \text{if } i \in N \setminus S. \end{cases}$$

Two subsets of agents, S and $N \setminus S$, can be connected to the source either separately or jointly. If there are no savings when they are jointly connected to the source, this property implies that agents will pay the same in both circumstances.

Notice that if a rule f satisfies PM, then also satisfies SEP.

Solidarity (SOL): For all *mcstp* (N_0, C) and (N_0, C') such that $C \leq C'$, we have

$$\psi(N_0, C) \leq \psi(N_0, C').$$

This property implies that if a number of connection costs increase and the rest of connection costs (if any) remain the same, no agent can be better off. Notice that *SOL* demands agents' contribution to move in the same direction irrespective of their locations on minimum cost spanning trees.

We say that two *mcstp* (N_0, C) and (N_0, C') are *tree-equivalent* if there exists a tree t such that, firstly, t is an *mt* for both (N_0, C) and (N_0, C') and secondly, $c_{ij} = c'_{ij}$ for all $(i, j) \in t$.

Independence of irrelevant trees (IIT): If two *mcstp* (N_0, C) and (N_0, C') are tree-equivalent,

$$f(N_0, C) = f(N_0, C').$$

IIT states that if two *mcstp* have the same *mt* with the same costs in each arc belonging to, then the rule must coincide in both *mcstp*.

Bergantiños and Vidal-Puga (2007a) prove that if a rule f satisfies SOL, then also satisfies IIT.

Equal sharing of extra costs (ESEC): Let (N_0, C) and (N_0, C') be two *mcstp*. Let $c_0, c'_0 \geq 0$. Assuming $c_{0i} = c_0$ and $c'_{0i} = c'_0$ for all $i \in N$, $c_0 < c'_0$, and $c_{ij} = c'_{ij} \leq c_0$ for all $i, j \in N$, we have

$$f_i(N_0, C') = f_i(N_0, C) + \frac{c'_0 - c_0}{n}$$

for all $i \in N$.

This property is interpreted as follows: a group of agents N faces a problem (N_0, C) in which all of them have the same connection cost to the source ($c_{i0} = c_0$) and in which this cost is greater than the connection costs between agents ($c_{ij} \leq c_0$). Under these circumstances, an optimal network implies that any one agent connects directly to the source, and that the rest connect to the source through this agent. Moreover, they agree that the correct solution is $f(N_0, C)$. Assume that an error was made and that the connection cost to the source is $c'_0 > c_0$. *ESEC* states that the agents should share this extra cost $c'_0 - c_0$ equally among them.

Bergantiños and Vidal-Puga (2007a) give the following characterizations of the folk solution φ .

Theorem 0:

- (a) φ is the unique rule on *mcstp* satisfying IIT, SEP, and ESEC.
- (b) φ is the unique rule on *mcstp* satisfying SOL, PM, and ESEC.

In Lemma 0 below we present some results used in the paper. The proof can be found in Bergantiños and Gómez-Rúa, 2009 and in Bergantiños and Vidal-Puga (2007a, 2007b, 2009).

Lemma 0. (a) Given (N_0, C, G) we can find an *mt* t in (N_0, C) satisfying:

(i) For each $k = 1, \dots, m$, t_{G^k} is an *mt* in (G^k, C) .

(ii) $t \setminus \left(\bigcup_{k=1}^m t_{G^k} \right)$ defined as $\{(k, k') : \exists i \in G^k, j \in G^{k'} \text{ with } (i, j) \in t\}$

is an *mt* in (G_0, C^G) .

(iii) For each $k = 1, \dots, m$ and each $i \in G^k$, $t_{G^k} \cup \{(0, i)\}$ is an *mt* in (G_0^k, C^φ) .

(b) If C is an irreducible matrix, then for all $S \subset N$ and $i \notin S$ we have that

$$v_C(S \cup \{i\}) - v_C(S) = \min_{j \in S_0} \{c_{ij}\}.$$

3 Minimum cost spanning tree problems with groups

In Bergantiños and Gómez-Rúa (2009) we introduce the *minimum cost spanning tree problems with groups*. An *mcstp* with groups is a triple (N_0, C, G) where (N_0, C) is an *mcstp*, $G = \{G^1, \dots, G^m\}$ is a partition of N and for each $k = 1, \dots, m$

$$\max_{i, j \in G^k} \{c_{ij}\} \leq \min_{i \in G^k, j \in N_0 \setminus G^k} \{c_{ij}\}.$$

A *rule* in *mcstp* with groups is a function f such that $f(N_0, C, G) \in \mathbb{R}^N$ and $\sum_{i \in N} f_i(N_0, C, G) = m(N_0, C)$ for each *mcstp* (N_0, C) .

As in classical *mcstp*, the main objective is to divide the cost associated with an *mt* among the agents in a fair way.

In Bergantiños and Gómez-Rúa (2009) we introduce a rule, F as the unique in *mcstp* with groups satisfying a set of properties. The rule is defined as follows: we first give the intuitive idea. This rule can be considered as a two step rule. In the first step, we compute the amount that each group should pay in order to be connected to the source. We do this by applying the rule φ defined in Bergantiños and Vidal-Puga (2007a).

In the second step we decide the amount that each agent of each group has to pay. For each group G^k , we consider the *mcstp* inside each group (G_0^k, C^φ) . In this *mcstp*, the connection cost between two agents of G^k is the same as in C but the connection cost between any agent of G^k and the source is the amount computed for G^k in the first step.

We now present the definition formally. Given the *mcstp* with groups (N_0, C, G) we define the *mcstp* among groups (G_0, C^G) as follows:

- $G_0 = \{G^0, G^1, \dots, G^m\}$ where $G^0 = 0$. In order to simplify the notation, often we use k instead of G^k .

- C^G is the cost matrix and for each $G^k, G^{k'} \in G_0$ the connection cost between G^k and $G^{k'}$ is denoted by

$$c_{kk'}^G = \min_{i \in G^k, j \in G^{k'}} \{c_{ij}\}.$$

Let (N_0, C, G) be an *mcstp* with groups. For each $i \in G^k$,

$$F_i(N_0, C, G) = \varphi_i(G_0^k, C^\varphi)$$

where

$$c_{jj'}^\varphi = \begin{cases} c_{jj'} & \text{if } 0 \notin \{j, j'\} \\ \varphi_k(G_0, C^G) & \text{if } 0 \in \{j, j'\}. \end{cases}$$

In this paper we follow the axiomatic approach and we present two axiomatic characterizations for the rule F . Our idea is to generalize the axiomatic characterizations of the folk rule φ given by Bergantiños and Vidal-Puga (2007a) and presented in Theorem 0.

In order to obtain these axiomatic approach, we now adapt these properties to *mcstp* with groups.

SOL and *IIT* can be formulated in a similar way. *PM*, *SEP*, and *CSEC* should be adapted. The main idea for adapting *PM* and *SEP* is claiming both twice, once among the groups and other among agents inside the same group. *CSEC* is adapted in a different way explained below.

Bergantiños and Gómez-Rúa (2009) introduce the following properties by adapting *PM*.

Population monotonicity over groups (PMG): For all *mcstp* with groups (N_0, C, G) , all $G^k \in G$, and all $i \in N \setminus G^k$,

$$f_i(N_0, C, G) \leq f_i((N \setminus G^k)_0, C, G \setminus G^k).$$

The population monotonicity over agents will show what happens when an agent enters in a group. We claim that no agent of the initial group can be worse off.

Assume that after the entrance of agent i in group G^k the minimum connection cost between group G^k and the rest of the groups did not change, *i.e.*, for each $G^l \in G_0 \setminus \{G^k\}$, $\min_{j \in G^k, j' \in G^l} \{c_{jj'}\} = \min_{j \in G^k \setminus \{i\}, j' \in G^l} \{c_{jj'}\}$. Since we are assuming that the amount paid by a group should not depend on the internal characteristics of the other groups, and the entrance of agent i do

not change the connection cost among groups, we claim that agents of the others groups must pay the same.

Population monotonicity over agents (PMA): For all *mcstp* with groups (N_0, C, G) , all $G^k \in G$, and all $i \in G^k$ such that $G^k \setminus \{i\} \neq \emptyset$,

$$f_j(N_0, C, G) \leq f_j((N \setminus \{i\})_0, C, G^{-i}) \text{ if } j \in G^k \setminus \{i\}.$$

Moreover, if for each $G^l \in G \setminus \{G^k\}$, $\min_{j \in G^k, j' \in G^l} \{c_{jj'}\} = \min_{j \in G^k \setminus \{i\}, j' \in G^l} \{c_{jj'}\}$, then

$$f_j(N_0, C, G) = f_j((N \setminus \{i\})_0, C, G^{-i}) \text{ if } j \in N \setminus G^k.$$

We now define two properties of separability.

Separability among groups (SEPG). For all *mcstp* (N_0, C, G) and $R \subset G$ satisfying $m(N_0, C) = m((\cup_{k \in R} G^k)_0, C) + m((\cup_{k \notin R} G^k)_0, C)$, we have

$$f_i(N_0, C, G) = \begin{cases} f_i\left(\left(\cup_{k \in R} G^k\right)_0, C, \left\{\left\{G^k\right\}\right\}_{k \in R}\right) & \text{if } i \in \cup_{k \in R} G^k \\ f_i\left(\left(\cup_{k \notin R} G^k\right)_0, C, \left\{\left\{G^k\right\}\right\}_{k \notin R}\right) & \text{if } i \in N \setminus \cup_{k \in R} G^k. \end{cases}$$

This property ensures the following: two subsets of groups R and $G \setminus R$ can be connected to the source either separately or jointly. If there is no savings when they are jointly connected to the source, this property implies that agents in each group will pay the same in both circumstances.

Separability among agents (SEPA). For all *mcstp* $(N_0, C, \{N\})$ and $S \subset N$ satisfying $m(N_0, C) = m(S_0, C) + m((N \setminus S)_0, C)$, we have

$$f_i(N_0, C, \{N\}) = \begin{cases} f_i(S_0, C, \{S\}) & \text{if } i \in S \\ f_i((N \setminus S)_0, C, \{N \setminus S\}) & \text{if } i \in N \setminus S. \end{cases}$$

Assume that there is a unique group. Two subsets of agents S and $N \setminus S$ can be connected to the source either separately or jointly. If there is no savings when they are jointly connected to the source, this property implies that agents will pay the same in both circumstances.

Finally, we adapt *ESEC* to *mcstp* with groups.

Coalitional sharing of extra costs (CSEC): Let (N_0, C, G) and (N_0, C', G) be two *mcstp* with groups. Let $c_0, c'_0 \geq 0$. Assume that $c_{0i} = c_0$ and $c'_{0i} = c'_0$ for all $i \in N$, $c_0 < c'_0$, and $c_{ij} = c'_{ij} \leq c_0$ for all $i, j \in N$. For each $i \in G^k \in G$

$$f_i(N_0, C', G) = f_i(N_0, C, G) + \frac{c'_0 - c_0}{m |G^k|}$$

CSEC states that when an extra cost appears, $c'_0 - c_0$, the groups should share this extra cost equally among them. Moreover, the extra cost assigned to each group should be shared equally among the agents of this group.

4 Characterization

In this section we generalize the results presented in Theorem 0 (Bergantiños and Vidal-Puga, 2007a) to *mcstp* with groups, using the properties presented in the previous section. We obtain two characterization results for the rule F introduced in Bergantiños and Gómez-Rúa (2009).

Theorem 1. (a) F is the unique rule satisfying IIT, SEPG, SEPA, and CSEC.

(b) F is the unique rule satisfying SOL, PMG, PMA, and CSEC.

Proof of Theorem 1. We first prove that F satisfies all the properties mentioned above. Bergantiños and Gómez-Rúa (2009) prove that F satisfies PMG and PMA. The other properties are a consequence of the following claims:

Claim 1. F satisfies SOL.

Proof of Claim 1. Let (N_0, C, G) and (N_0, C', G) such that $C \leq C'$. Consider the problems among coalitions, (G_0, C^G) and (G_0, C'^G) . By definition of C^G and C'^G , we have that $c_{kk'}^G \leq c_{kk'}'^G$ for all $G^k, G^{k'} \in G_0$. Then $C^G \leq C'^G$. Since φ satisfies *SOL* (Theorem 0), $\varphi_k(G_0, C^G) \leq \varphi_k(G_0, C'^G)$ for all $G^k \in G_0$.

Consider now the problem inside a coalition (G_0^k, C'^φ) , $k = 1, \dots, m$. In this case, $c_{ij}^\varphi = c_{ij} \leq c'_{ij} = c_{ij}'^\varphi$ for all $i, j \in G^k$ and $c_{0i}^\varphi = \varphi_k(G_0, C^G) \leq \varphi_k(G_0, C'^G) = c_{0i}'^\varphi$ for all $i \in G^k$. Then, $C^\varphi \leq C'^\varphi$, since φ satisfies *SOL* (Theorem 0), we have

$$F_i(N_0, C, G) = \varphi_i(G_0^k, C^\varphi) \leq \varphi_i(G_0^k, C'^\varphi) = F_i(N_0, C', G)$$

for all $i \in G^k$. ■

Claim 2. Let f be a rule satisfying *PMG*, then f also satisfies *SEPG*.

Proof of Claim 2. Let f be a rule satisfying *PMG* and $R \subset G$ as in the definition of *SEPG*. Under *PMG* we know that $f_i(N_0, C, G) \leq$

$f_i \left((\cup_{k \in R} G^k)_0, C, \{\{G^k\}\}_{k \in R} \right)$ for all $i \in \cup_{k \in R} G^k$ and $f_i(N_0, C, G) \leq f_i \left((N \setminus \cup_{k \in R} G^k)_0, C, \{\{G^k\}\}_{k \in R} \right)$ for all $i \in N \setminus \cup_{k \in R} G^k$.

Moreover by definition of f ,

$$\begin{aligned} \sum_{i \in N} f_i(N_0, C, G) &= m(N_0, C), \\ \sum_{i \in \cup_{k \in R} G^k} f_i \left((\cup_{k \in R} G^k)_0, C, \{\{G^k\}\}_{k \in R} \right) &= m((\cup_{k \in R} G^k)_0, C), \text{ and} \\ \sum_{i \in (N \setminus \cup_{k \in R} G^k)} f_i \left((N \setminus \cup_{k \in R} G^k)_0, C, \{\{G^k\}\}_{k \in R} \right) &= m((N \setminus \cup_{k \in R} G^k)_0, C). \end{aligned}$$

Since $m(N_0, C) = m((\cup_{k \in R} G^k)_0, C) + m((N \setminus \cup_{k \in R} G^k)_0, C)$, it is clear that f satisfies SEPG. ■

Claim 3. Let f be a rule satisfying *PMA*, then f also satisfies *SEPA*.

Proof of Claim 3. It is similar to the proof of Claim 2 and we omit it. ■

Claim 4. F satisfies CSEC.

Proof of Claim 5. Let (N_0, C, G) and (N_0, C', G) be under the conditions of CSEC. Consider the problems among groups (G_0, C^G) and (G_0, C'^G) . Thus, $c_{0k}^G = \min_{j \in G^k} \{c_{0j}\} = c_0$ and $c_{0k}^{G'} = c'_0$ for all $G^k \in G$. Moreover, $c_{kk'}^G = \min_{j \in G^k, j' \in G^{k'}} \{c_{jj'}\} \leq c_0$, $c_{kk'}^{G'} = \min_{j \in G^k, j' \in G^{k'}} \{c'_{jj'}\} \leq c'_0$, and $c_{kk'}^G \leq c_{kk'}^{G'}$ for all $G^k, G^{k'} \in G$. Reasoning in the same way, we obtain that $c_{0k}^{G'} = c'_0$ and $c_{kk'}^{G'} = c_{kk'}^G$. Thus, the problems (G_0, C^G) and (G_0, C'^G) are under the conditions of ESEC. Since φ satisfies ESEC (Theorem 0), we have that

$$\varphi_k(G_0, C'^G) = \varphi_k(G_0, C^G) + \frac{c'_0 - c_0}{m} \quad (1)$$

for all $G^k \in G$.

Consider now the problems inside each group $G^k \in G$, (G_0^k, C^φ) and (G_0^k, C'^φ) . By definition, $c_{0i}^\varphi = \varphi_k(G_0, C^G) := c_0^\varphi$ for all $i \in G^k$ and $c_{ij}^\varphi = c_{ij}$ for all $i, j \in G^k$. Moreover, in the proof of Lemma 0 (a) (iii) (Bergantiños and Gómez-Rúa, 2009) we prove that $c_0^\varphi \geq \max_{j, j' \in G^k} \{c_{jj'}^\varphi\}$. Reasoning in the same way we obtain that $c_{0i}^{I\varphi} = \varphi_k(G_0, C'^G) = c_0^{I\varphi}$ for all $i \in G^k$, $c_{ij}^{I\varphi} = c_{ij}^I = c_{ij}$ for

all $i, j \in G^k$ and $c_0^{i\varphi} \geq \max_{j, j' \in G^k} \{c_{jj'}^{i\varphi}\}$. Furthermore, taking into account expression (1), we have that $c_0^\varphi < c_0^{i\varphi}$, and $c_0^{i\varphi} - c_0^\varphi = \varphi_k(G_0, C^{iG}) - \varphi_k(G_0, C^G) = \frac{c_0^{i\varphi} - c_0^\varphi}{m}$. Thus, (G_0^k, C^φ) and $(G_0^k, C^{i\varphi})$ are also under the conditions of ESEC. Since φ satisfies ESEC (Theorem 0), we obtain that

$$\begin{aligned} F_i(N_0, C', G) &= \varphi_i(G_0^k, C^{i\varphi}) = \varphi_i(G_0^k, C^\varphi) + \frac{c_0^{i\varphi} - c_0^\varphi}{|G^k|} \\ &= \varphi_i(G_0^k, C^\varphi) + \frac{c_0^{i\varphi} - c_0^\varphi}{m |G^k|} \\ &= F_i(N_0, C, G) + \frac{c_0^{i\varphi} - c_0^\varphi}{m |G^k|} \end{aligned}$$

for all $i \in G^k$. ■

We now prove the uniqueness in each of the two parts.

(a) We apply an induction argument over the number of agents $|N|$. If $|N| = 1$ the result is trivial. Assume that the result holds for less than n agents. We prove it for n agents.

Let f be a rule satisfying these properties. Given an *mcstp* with groups (N_0, C, G) we prove that $f(N_0, C, G) = F(N_0, C, G)$.

Let $t = \{(i^0, i)\}_{i \in N}$ be an *mt* in (N_0, C) as in Lemma 0 (a).

Assume that there is at least two agents who connect directly to the source in t . Namely, $|\{i \in N : (0, i) \in t\}| \geq 2$. Let j, j' be such that $(0, j) \in t$ and $(0, j') \in t$,

We define $S = \text{Fol}(j, t) \cup \{j'\}$. Thus, $m(N_0, C) = m(S_0, C) + m((N \setminus S)_0, C)$. Since t satisfies the conditions of Lemma 0 (a), there exists $R \subset \{1, \dots, m\}$ such that $S = \cup_{k \in R} G^k$. Since f satisfies SEPG, we have that

$$f_i(N_0, C, G) = \begin{cases} f_i\left(\left(\cup_{k \in R} G^k\right)_0, C, \{\{G^k\}\}_{k \in R}\right) & \text{if } i \in \cup_{k \in R} G^k \\ f_i\left(\left(\cup_{k \notin R} G^k\right)_0, C, \{\{G^k\}\}_{k \notin R}\right) & \text{if } i \in N \setminus \cup_{k \in R} G^k. \end{cases}$$

Since $j \in \cup_{k \in R} G^k$ and $j' \in \cup_{k \notin R} G^k$, by the induction hypothesis,

$$\begin{aligned} f_j\left(\left(\cup_{k \in R} G^k\right)_0, C, \{\{G^k\}\}_{k \in R}\right) &= F_j\left(\left(\cup_{k \in R} G^k\right)_0, C, \{\{G^k\}\}_{k \in R}\right) \text{ and} \\ f_{j'}\left(\left(\cup_{k \notin R} G^k\right)_0, C, \{\{G^k\}\}_{k \notin R}\right) &= F_{j'}\left(\left(\cup_{k \notin R} G^k\right)_0, C, \{\{G^k\}\}_{k \notin R}\right). \end{aligned}$$

Since F also satisfies SEPG, we conclude that

$$\begin{aligned} F_i \left(\left(\bigcup_{k \in R} G^k \right)_0, C, \left\{ \{G^k\} \right\}_{k \in R} \right) &= F_i(N_0, C, G) \text{ if } i \in \bigcup_{k \in R} G^k \text{ and} \\ F_i \left(\left(\bigcup_{k \notin R} G^k \right)_0, C, \left\{ \{G^k\} \right\}_{k \notin R} \right) &= F_i(N_0, C, G) \text{ if } i \in N \setminus \bigcup_{k \in R} G^k. \end{aligned}$$

Thus, $f_i(N_0, C, G) = F_i(N_0, C, G)$ for all $i \in N$.

Assume now that there is a unique agent who connect directly to the source in t . We define $x = \max \{c_{i^0 i} : (i^0, i) \in t\}$ and $x' = \max \{c_{i^0 i} : (i^0, i) \in t \text{ and } c_{i^0 i} < x\}$.

We consider several cases.

1. There exists $(j^0, j) \in t$ such that $j \in G^k, j^0 \in G^{k'}, k \neq k'$, and $c_{j^0 j} = x$.

By the definition of C^* as the minimal network associated with t , we can deduce that t is also an mt in C^* . Since (j^0, j) is in the unique path in t from j to 0, we deduce that $c_{0j}^* = x$.

Since f and F satisfy *IIT*, $f(N_0, C, G) = f(N_0, C^*, G)$ and $F(N_0, C, G) = F(N_0, C^*, G)$. So we can focus in the problem (N_0, C^*, G) .

Let $t^* = t \setminus \{(j^0, j)\} \cup \{(0, j)\}$. It is clear that $c(C^*, t^*) = c(C^*, t)$.

Then, t^* is also an mt in C^* . Moreover, there are at least two agents connected directly to the source in t^* . Thus, $f(N_0, C^*, G) = F(N_0, C^*, G)$.

2. We are not in Case 1 and there exists $(j^0, j) \in t$ such that $j, j^0 \in G^k$ and $c_{j^0 j} = x$.

We first prove that $G = \{N\}$. Suppose not. By definition of C , there exists an arc $(i^0, i) \in t$ with $i \in G^k, i^0 \in N_0 \setminus G^k$ such that $c_{i^0 i} \geq c_{j^0 j}$. Since $c_{j^0 j} = x, c_{i^0 i} = c_{j^0 j} = x$. So we are in Case 1, which is a contradiction.

We consider C^* . Using arguments similar to those used in Case 1, SEPA instead SEPG, we can deduce that $f(N_0, C, G) = F(N_0, C, G)$.

3. There exists $j \in N$ such that $(0, j) \in t$ and $c_{0j} = x > c_{i^0 i}$ for all $(i^0, i) \in t$ with $\{i^0, i\} \subset N$. Notice that we are not in any of the previous cases.

We can compute C^* as the minimal network associated to t . Since $(0, j)$ is the unique arc in t containing the source, for each $i \in N, (0, j)$ is in the unique path in t from 0 to i . Thus, $c_{0i}^* = c_{0j} = x$. Moreover, for

each $\{i, i'\} \subset N$, $(0, j)$ is not in the unique path in t from i to i' . Thus, $c_{ii'}^* \leq x' < x$.

Since f and F satisfy *IIT*, $f(N_0, C, G) = f(N_0, C^*, G)$ and $F(N_0, C, G) = F(N_0, C^*, G)$.

Let C' be such that $c'_{0i} = x'$ for all $i \in N$ and $c'_{ih} = c_{ih}^*$ otherwise. Applying Prim's algorithm is easy to see that t is also an mt in C' . Moreover C^* and C' are under the hypothesis of *CSEC*. Since f and F satisfy *CSEC*, it is enough to prove that $f(N_0, C', G) = F(N_0, C', G)$.

Let $(k^0, k) \in t$ be such that $c_{k^0k} = x'$. Thus, $c_{k^0k}^* = c'_{k^0k} = x'$. Let $t' = t \setminus \{(k^0, k)\} \cup \{(0, k)\}$. It is clear that $c(C', t') = c(C', t)$. Then, t' is also an mt in C' .

Two cases are possible. First, we can take $k^0 \in G^l$ and $k \in N \setminus G^l$. Then, t' satisfies the conditions of Lemma 0 (a). Moreover, there are at least two agents connected directly to the source in t' . Thus, $f(N_0, C', G) = F(N_0, C', G)$. Second, for any arc $(k^0, k) \in t$ with $c_{k^0k} = x'$ we have that $k, k^0 \in G^l$. Using arguments similar to those used in Case 2 we can deduce that $G = \{N\}$ and $f(N_0, C', G) = F(N_0, C', G)$. ■

(b) Since *SOL* implies *IIT*, *PMG* implies *SEPG*, and *PMA* implies *SEPA*, it is a trivial consequence of part (a). ■

Proposition 1. The properties used in Theorem 1 (a) and (b) are independent.

Proof: We proof that if we remove some of the properties of Theorem 1 (a) and (b), we can find more rules satisfying the other properties. We do this by considering several claims.

Claim 1 There exist rules satisfying *SOL* (and hence *IIT*), *PMG* (and hence *SEPG*), *PMA* (and hence *SEPA*), and failing *CSEC*.

Proof of Claim 1: We define the rule f^1 as follows. Given $T \subset \mathcal{N}$, let π^N denote the order induced in N by the index of the agents. Namely, given $i, j \in N$, $\pi^N(i) < \pi^N(j)$ if and only if $i < j$. For each $mcstp(N_0, C)$ and $i \in N$ we define

$$\psi_i(N_0, C) = v_{C^*}(Pre(i, \pi^N) \cup \{i\}) - v_{C^*}(Pre(i, \pi^N)).$$

Let (N_0, C, G) be an *mcstp* with groups and $i \in G^k$. Thus,

$$f_i^1(N_0, C, G) = \psi_i(G_0^k, C^\varphi).$$

1. f^1 satisfies SOL. Let (N_0, C, G) and (N_0, C', G) be such that $C \leq C'$. Consider the problems among coalitions, (G_0, C^G) and (G_0, C'^G) . Then, $C^G \leq C'^G$. Since φ satisfies SOL (Theorem 0), $\varphi_k(G_0, C^G) \leq \varphi_k(G_0, C'^G)$ for all $G^k \in G_0$.

Consider now the problem inside a coalition G_0^k with $k = 1, \dots, m$. Then, $C^\varphi \leq C'^\varphi$. By definition of the irreducible matrix, $(C^\varphi)^* \leq (C'^\varphi)^*$. By Lemma 0 (b), for all $i \in G^k$,

$$\psi_i(G_0^k, C^\varphi) \leq \psi_i(G_0^k, C'^\varphi).$$

2. f^1 satisfies PMG. See Bergantiños and Gómez-Rúa (2009).
3. f^1 satisfies PMA. See Bergantiños and Gómez-Rúa (2009).
4. f^1 fails CSEC. Consider the *mcstp* with groups where $N = \{1, 2\}$, $G = \{\{i\}\}_{i \in N}$, and matrices

$$C = \begin{pmatrix} 0 & 10 & 10 \\ 10 & 0 & 2 \\ 10 & 2 & 0 \end{pmatrix} \text{ and } C' = \begin{pmatrix} 0 & 12 & 12 \\ 12 & 0 & 2 \\ 12 & 2 & 0 \end{pmatrix}.$$

In this case, $f^1(N_0, C, G) = (10, 2)$ and $f^1(N_0, C', G) = (12, 2)$. ■

Claim 2 There exist rules satisfying SOL (and hence IIT), PMG (and hence SEPG), CSEC and failing SEPA (and hence PMA).

Proof of Claim 2: We define the rule f^2 as follows. For each *mcstp* (N_0, C) and $i \in N$ we define the equal division rule as

$$E_i(N_0, C) = \frac{m(N_0, C)}{|N|}.$$

Let (N_0, C, G) be an *mcstp* with groups and $i \in G^k$. Thus,

$$f_i^2(N_0, C, G) = E_i(G_0^k, C^\varphi).$$

1. f^2 satisfies SOL. Let (N_0, C, G) and (N_0, C', G) be such that $C \leq C'$. Then $C^G \leq C'^G$. Since φ satisfies SOL (Theorem 0), $\varphi_k(G_0, C^G) \leq \varphi_k(G_0, C'^G)$ for all $G^k \in G_0$.

Consider now the problem inside a coalition (G_0^k, C^φ) with $k = 1, \dots, m$. Then, $C^\varphi \leq C'^\varphi$ and hence

$$f_i^2(N_0, C, G) = E_i(G_0^k, C^\varphi) \leq E_i(G_0^k, C'^\varphi) = f_i^2(N_0, C', G)$$

for all $i \in G^k$.

2. f^2 satisfies *PMG*. Let $G^k \in G$. Since φ satisfies *PM* (Theorem 0), $\varphi_l(G_0, C^G) \leq \varphi_l((G \setminus G^k)_0, C^{G \setminus G^k})$ for all $l \neq k$. Moreover, $((G \setminus G^k)_0, C^G) = ((G \setminus G^k)_0, C^{G \setminus G^k})$

Let C'^φ denote the matrix C^φ associated with the problem $((N \setminus G^k)_0, C, G \setminus G^k)$. Let $G^l \in G \setminus G^k$. For all $i, j \in G^l$, $c_{ij}^\varphi = c_{ij}'^\varphi$. For all $i \in G^l$,

$$c_{0i}^\varphi = \varphi_l(G_0, C^G) \leq \varphi_l((G \setminus G^k)_0, C^{G \setminus G^k}) = c_{0i}'^\varphi.$$

That is $C^\varphi \leq C'^\varphi$. Then, $E_i(G_0^l, C^\varphi) \leq E_i(G_0^l, C'^\varphi)$ and hence,

$$f_i^2(N_0, C, G) \leq f_i^2((N \setminus G^k)_0, C, G \setminus G^k).$$

3. f^2 satisfies *CSEC*. The proof is similar than the one for F and we omit it.

4. f^2 fails *SEPA*. Consider the *mcstp* with groups where $N = \{1, 2, 3\}$, $G = \{N\}$, and matrix

$$C = \begin{pmatrix} 0 & 1 & 2 & 1 \\ 1 & 0 & 0 & 1 \\ 2 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix}$$

In this case, it is clear that $m(N_0, C) = m(\{1, 2\}_0, C) + m(\{3\}_0, C)$. However, $f^2(N_0, C, G) = (\frac{2}{3}, \frac{2}{3}, \frac{2}{3})$ and $f^2(\{1, 2\}_0, C) = (\frac{1}{2}, \frac{1}{2})$ and $f^2(\{3\}_0, C) = 1$. ■

Claim 3 There exist rules satisfying *SOL* (and hence *IIT*), *PMA* (and hence *SEPA*), *CSEC* and failing *SEPG* (and hence *PMG*).

Proof of Claim 3: We define the rule f^3 as follows. Let (N_0, C, G) be an *mcstp* with groups and $i \in G^k$. Thus,

$$f_i^3(N_0, C, G) = \varphi_i(G_0^k, C^E)$$

where $c_{jj'}^E = \begin{cases} c_{jj'} & \text{if } 0 \notin \{j, j'\} \\ E_k(G_0, C^G) & \text{if } 0 \in \{j, j'\}. \end{cases}$

1. f^3 satisfies *SOL*. Let (N_0, C, G) and (N_0, C', G) such that $C \leq C'$. Then $C^G \leq C'^G$. Hence, $E_k(G_0, C^G) \leq E_k(G_0, C'^G)$ for all $G^k \in G_0$.

Consider now the problem inside a coalition (G_0^k, C^E) with $k = 1, \dots, m$. Then, $C^E \leq C'^E$. Since φ satisfies *SOL* (Theorem 0),

$$f_i^3(N_0, C, G) = \varphi_i(G_0^k, C^E) \leq \varphi_i(G_0^k, C'^E) = f_i^3(N_0, C', G)$$

for all $i \in G^k$.

2. f^3 satisfies *PMA*. Let $G^k \in G$ and $i \in G^k$. By convenience, let us denote as C' the cost matrix C restricted to the problem $((N \setminus \{i\})_0, C, G^{-i})$. Notice that C' coincides with C on the agents of $(N \setminus \{i\})_0$.

We consider several cases:

a) Assume that $c_{kl}^G = c_{kl}'^G$ for all $l \in \{0, 1, \dots, m\}$. Thus, $E_l(G_0, C^G) = E_l(G^{-i}, C'^{G^{-i}})$ for all $l = 1, \dots, m$.

- Let $j \in G^k \setminus \{i\}$. Since φ satisfies *PM* (Theorem 0), $\varphi_j(G_0^k, C^E) \leq \varphi_j((G^k \setminus \{i\})_0, C^E)$. Then,

$$\begin{aligned} f_j^3(N_0, C, G) &= \varphi_j(G_0^k, C^E) \leq \varphi_j((G^k \setminus \{i\})_0, C^E) \\ &= \varphi_j((G^k \setminus \{i\})_0, C'^E) \\ &= f_j^3((N \setminus \{i\})_0, C', G^{-i}). \end{aligned}$$

- Let $G^l \in G$ such that $l \neq k$. Then, $c_{jj'}^E = c_{jj'}'^E$ for all $j, j' \in G^l \cup \{0\}$. Hence,

$$f_j^3(N_0, C, G) = f_j^3((N \setminus \{i\})_0, C', G^{-i}).$$

b) Assume that $c_{kk'}^G \neq c_{kk'}'^G$ for some $k' \in \{0, 1, \dots, m\}$. Then, $c_{kk'}^G < c_{kk'}'^G$. Moreover, $c_{ll'}^G \leq c_{ll'}'^G$ for all $l, l' \in \{0, 1, \dots, m\}$.

Then, $E_k(G_0, C^G) \leq E_k(G_0, C'^G)$, $c_{jj'}^E = c_{jj'}'^E$ for all $j, j' \in G^k \setminus \{i\}$, and $c_{0j}^E \leq c_{0j}'^E$ for all $j \in G^k \setminus \{i\}$. Since φ satisfies *SOL* (Theorem 0), $\varphi_j((G^k \setminus \{i\})_0, C^E) \leq \varphi_j((G^k \setminus \{i\})_0, C'^E)$ for all $j \in G^k \setminus \{i\}$.

Since φ satisfies *PM* (Theorem 0), $\varphi_j(G_0^k, C^E) \leq \varphi_j((G^k \setminus \{i\})_0, C^E)$. Then, for all $j \in G^k \setminus \{i\}$

$$\begin{aligned} f_j^3(N_0, C, G) &= \varphi_j(G_0^k, C^E) \leq \varphi_j((G^k \setminus \{i\})_0, C^E) \\ &\leq \varphi_j((G^k \setminus \{i\})_0, C'^E) \\ &= f_j^3((N \setminus \{i\})_0, C', G^{-i}). \end{aligned}$$

3. f^3 satisfies *CSEC*. The proof is similar than the one for F and we omit it.

4. f^3 fails *SEPG*. Consider the *mcstp* with groups where $N = \{1, 2, 3\}$, $G = \{\{i\}\}_{i \in N}$, and matrix

$$C = \begin{pmatrix} 0 & 1 & 2 & 1 \\ 1 & 0 & 0 & 1 \\ 2 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix}$$

In this case, it is clear that $m(N_0, C) = m(\{1, 2\}_0, C) + m(\{3\}_0, C)$. However, $f^3(N_0, C, G) = (\frac{2}{3}, \frac{2}{3}, \frac{2}{3})$ and $f^3(\{1, 2\}_0, C) = (\frac{1}{2}, \frac{1}{2})$ and $f^3(\{3\}_0, C) = 1$. ■

Claim 4 There exist rules satisfying *PMG* (and hence *SEPG*), *PMA* (and hence *SEPA*), *CSEC* and failing *SOL* (and hence *IIT*).

Proof of Claim 4: We define the rule f^4 as follows. Let (N_0, C, G) be an *mcstp* with groups. We say that $\pi \in \Pi_N$ is *admissible* if the following conditions are satisfied: i) given $i \in G^k$, $j \in G^{k'}$, $k \neq k'$, if $\pi(i) < \pi(j)$, then $c_{0k}^G \leq c_{0k'}^G$; ii) given $i, j \in G^k$, if $\pi(i) < \pi(j)$, then $c_{0i} \leq c_{0j}$ and iii) if $\pi(i) < \pi(j) < \pi(l)$, with $i, l \in G^k$, then, $j \in G^k$. We denote by Π'^N the set of all permutations over N that are admissible. In an intuitive way: we first order the groups by decreasing connection cost to the source; and then the agents inside groups following the same idea.

Thus, for all $i \in N$,

$$f_i^4(N_0, C, G) = \frac{1}{|\Pi'_N|} \sum_{\pi \in \Pi'_N} v_{C^*}(Pre(i, \pi) \cup \{i\}) - v_{C^*}(Pre(i, \pi)).$$

1. f^4 satisfies *PMG*. Making some computations we can prove it. Even the proof is not trivial, we do not think that it is specially relevant and we omit it.

2. f^4 satisfies *PMA*. Making some computations we can prove it. Even the proof is not trivial, we do not think that it is specially relevant and we omit it.

3. f^4 satisfies *CSEC*. Let (N_0, C, G) and (N_0, C', G) be two *mcstp* with groups under the conditions of *CSEC*. Then, a permutation $\pi \in \Pi'^N$ is admissible if given $i, i' \in G^k \in G$ and $j \in N$ with $\pi(i) < \pi(j) < \pi(i')$, then $j \in G^k$. Therefore, $f^4(N_0, C, G) = Ow(N, v_{C^*}, G)$. In Bergantiños and

Gómez-Rúa (2009) we prove that $F(N_0, C, G) = Ow(N, v_{C^*}, G)$. Then, in this case, $f^4(N_0, C, G) = F(N_0, C, G)$, and we have proved above that F satisfies *CSEC*.

4. f^4 fails *SOL*. Consider the *mcstp* with groups (N_0, C, G) and (N_0, C', G) where $N = \{1, 2\}$, $G = \{\{i\}\}_{i \in N}$, and matrices

$$C = \begin{pmatrix} 0 & 10 & 20 \\ 10 & 0 & 0 \\ 20 & 0 & 0 \end{pmatrix} \text{ and } C' = \begin{pmatrix} 0 & 30 & 20 \\ 30 & 0 & 0 \\ 20 & 0 & 0 \end{pmatrix}.$$

In this case, $f^4(N_0, C, G) = (10, 0)$ and $f^4(N_0, C', G) = (0, 20)$. ■

5 Acknowledgements

Financial support from Ministerio de Ciencia y Tecnología and FEDER through grant ECO2008-03484-C02-01/ECON and from Xunta de Galicia through grants PGIDIT06PXIB362390PR and INCITE08PXIB300005PR is gratefully acknowledged.

6 References

- Aarts, H. and Driessen, T. (1993) *The irreducible core of a minimum cost spanning tree game*. *Mathematical Methods of Operations Research* 38, 163-174.
- Bergantiños, G. and Gómez-Rúa, M. (2009) *Minimum cost spanning tree problems with groups*. Forthcoming in *Economic Theory*.
- Bergantiños, G. and Lorenzo, L. (2004) *A non-cooperative approach to the cost spanning tree problem*. *Mathematical Methods of Operations Research* 59, 393-403.
- Bergantiños, G. and Lorenzo, L. (2005) *Optimal equilibria in the non-cooperative game associated with cost spanning tree problems*. *Annals of Operations Research* 137, 101-115.
- Bergantiños, G. and Lorenzo, L. (2008) *Non cooperative cost spanning tree problems with budget restrictions*. *Naval Research Logistics* 55: 747-757.
- Bergantiños, G., Lorenzo, L., and Lorenzo-Freire, S. (2009). *The family of cost monotonic and cost additive rules in minimum cost spanning tree problems*. Forthcoming in *Social Choice and Welfare*.

Bergantiños, G. and Lorenzo-Freire, S. (2008) *Optimistic weighted Shapley rules in minimum cost spanning tree problems*. European Journal of Operational Research 185, 289-298.

Bergantiños, G. and Vidal-Puga, J.J. (2007a) *A fair rule in minimum cost spanning tree problems*. Journal of Economic Theory 137, 326-352.

Bergantiños, G. and Vidal-Puga, J.J. (2007b) *The optimistic TU game in minimum cost spanning tree problems*. International Journal of Game Theory 36, 223-239.

Bergantiños, G. and Vidal-Puga, J.J. (2009) *Additivity in minimum cost spanning tree problems*. Journal of Mathematical Economics 45, 38-42.

Bird, C. G., (1976) *On cost allocation for a spanning tree: A game theoretic approach*. Networks 6, 335-350.

Bogomolnaia, A., Moulin, H. (2008). *Sharing the cost of a minimal cost spanning tree: beyond the folk solution*. Mimeo, Rice University. Available at <http://www.ruf.rice.edu/~econ/faculty/moulin.html>

Branzei, R., Moretti, S., Norde, H. and Tijs, S. (2004) *The P-value for cost sharing in minimum cost spanning tree situations*. Theory and Decision 56, 47-61.

Chun, Y. and Lee, J. (2009). *Sequential contributions rules for minimum cost spanning tree problems*. Mimeo, Seoul University.

Dutta, B. and Kar, A. (2004) *Cost monotonicity, consistency and minimum cost spanning tree games*. Games and Economic Behavior 48, 223-248.

Feltkamp, V., Tijs, S. and Muto, S. (1994) *On the irreducible core and the equal remaining obligation rule of minimum cost extension problems*. Mimeo, Tilburg University.

Kar A., (2002) *Axiomatization of the Shapley value on minimum cost spanning tree games*. Games and Economic Behavior 38, 265-277.

Owen, G. (1977) *Values of games with a priori unions*. In: Henn R., O. Moeschlin, O., (Eds.), *Essays in Mathematical Economics and Game Theory*. Springer-Verlag, New York, pp. 76-88.

Prim, R.C. (1957) *Shortest connection networks and some generalizations*. Bell Systems Technology Journal 36, 1389-1401.

Shapley, L. S., (1953) *A value for n-person games*. In: H.W. Kuhn and A.W. Tucker, eds., *Contributions to the Theory of Games II* (Princeton, NJ) 307-317.